



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/00	A2	(11) International Publication Number: WO 98/58633 (43) International Publication Date: 30 December 1998 (30.12.98)
(21) International Application Number: PCT/CA98/00603 (22) International Filing Date: 22 June 1998 (22.06.98) (30) Priority Data: 60/050,491 23 June 1997 (23.06.97) US 60/086,750 27 May 1998 (27.05.98) US (71) Applicant (for all designated States except US): QUEEN'S UNIVERSITY AT KINGSTON [CA/CA]; Kingston, Ontario K7L 3N6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): ADAMS, Michael, A. [CA/CA]; 341 Renda Street, Kingston, Ontario K7M 5X9 (CA). HEATON, Jeremy, P., W. [CA/CA]; R.R. #1, Kingston, Ontario K7L 4V1 (CA). BANTING, James, D. [CA/CA]; Unit 1, Apartment 201, 47 Van Order Drive, Kingston, Ontario K7M 1B6 (CA). (74) Agent: STEEG, Carol, Miernicki; Parteq Innovations, Queen's University, Kingston, Ontario K7L 3N6 (CA).	(81) Designated States: AU, BR, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: MICRODOSE THERAPY (57) Abstract <p>Methods for treating vascular conditions associated with localized imbalance in vascular tone, which are hypothesized to be largely due to elevated endothelin (ET), are provided. The methods involve administration of nitric oxide (NO), agents which are able to provide NO, such as NO donors, agents which activate guanyl cyclase, such as YC-1, or agents which prolong the actions of endogenous NO or cyclic guanosine monophosphate (cGMP; a 2nd messenger molecule), such as phosphodiesterase (PDE) inhibitors. According to the invention, such agents are administered in minimal doses or microdoses by any route known in the art, so as to provide dosages which are about one half to about one twentieth (1/2 to 1/20) of those known to induce vasodilation in "normal" circulations. The low doses of these agents effectively alleviate vascular conditions associated with a reduction in NO production or an attenuation of NO effect, by restoring balance in vascular tone while exerting almost no systemic effect in normal vasculature.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

MICRODOSE THERAPY

Related Applications

This application claims the benefit of priority of U.S. Provisional Patent Applications No. 60/050,491, filed June 23, 1997, and No. 60/086,750, filed May 27, 1998. The contents of both these provisional applications are incorporated herein by reference.

Field of the Invention

The field of invention is the treatment of conditions concerned with peripheral vasoconstriction. More particularly, the invention is concerned with methods for establishing normal vascular tone in regions of the circulation which demonstrate pathophysiology. In particular, the invention concerns a method of treating erectile dysfunction by provision of nitric oxide, nitric oxide producing agents, or activators of guanyl cyclase in small doses or microdoses, i.e., doses that do not induce undesirable side effects, such as systemic vasodilation, under normal conditions.

Background of the Invention

It is widely known that administration of nitric oxide (NO), or compounds which deliver NO (i.e., NO donors, NO producing agents) to a subject, can provoke powerful vasodilator responses. Such administration is often accompanied by a number of undesirable side effects which include headache, flushing, and hypotension.

The physiological role of NO has been described as that of a powerful chronic vasodilator agent based on there being a marked increase in vascular tone following NO synthase (NOS) blockade (Johnson *et al.*, *Am. J. Hypertens.* 5:919, 1992; Tolins *et al.*, *Hypertens.* 17:909, 1991). The role of NO as a chronic vasodilator has only been inferred by indirect means, i.e., by removal of NOS activity. Endogenously, much more multiplicity and overlap in the control of vasodilation can be inferred from

the scientific literature. For example, vasodilation can be induced by acetylcholine, bradykinin, adenosine, adenosine triphosphate (ATP), histamine, vasoactive intestinal polypeptide (VIP), and leukotrienes, amongst others. The actions of these endogenous modulators have been shown to be dependent, at least in part, on the presence of the endothelium, an effect likely mediated by endothelial derived relaxing factor/NO (EDRF/NO) (Garg, U.C. *et al.*, *J. Biol. Chem.* 266:9, 1991; Garg, U.C. *et al.*, *J. Clin. Invest.* 83:1774, 1989; Palmer, R.M.J. *et al.*, *Nature* 327:524, 1987). Other vasodilator mechanisms exist which are not endothelium dependent, such as β_2 -adrenergic receptor activation, atrial natriuretic peptide (ANP) and certain prostaglandins. The actions of NO have been suggested to be mostly cGMP-mediated via guanylate cyclase activation, although other mechanisms have been suggested. For example, Garg *et al.* (*J. Biol. Chem.* 266:9, 1991; *J. Clin. Invest.* 83:1774, 1989) and others (Assender, J.W. *et al.*, *J. Cardiovasc. Pharmacol.* 17(Suppl.3):S104, 1991; O'Conner, K.J. *et al.*, *J. Cardiovasc. Pharmacol.* 17(Suppl.3):S100-S103, 1991) demonstrated a difference in the effects of NO-generating vasodilator agents in inhibiting vascular smooth muscle cell growth in culture; however, it is clear that NO can act not only as a vasodilator but also to inhibit vascular growth responses in a number of conditions (Farhy, R.D. *et al.*, *Circ. Res.* 72:1202, 1993).

It has been believed and widely practised that NO, in humans and animals, produced via sodium nitroprusside (SNP) infusion, causes vasodilation in peripheral vasculature at doses greater than 10 μ g/kg per min. It has recently been determined that NO also performs a function through interaction with endothelin (ET) (Banting *et al.*, *J. Hypertens.* 14:975, 1996; Richard *et al.*, *Circulation* 91:771, 1995). Prior to this time, ET had been believed to play a minimal role in maintaining tone in the peripheral microvasculature and to have little impact on the state of contraction of smooth muscle in those vessels. Recent studies have indicated (Banting *et al.*, *J. Hypertens.* 14:975, 1996) that ET is under the inhibitory control of NO and that administration of NOS inhibitors results in elevated levels of ET.

Endothelins were first described in 1988 and have been shown to be powerful vasoconstrictors, predominantly found in vascular endothelium and, since that time, numerous ET antagonists and their pharmaceutically acceptable salts have been identified and can be obtained commercially (e.g., Sigma, American Peptides).

5 Detailed descriptions of the chemical structures of various ET antagonists may be found in U.S. Patents No. 5,284,828 issued 8 February 1994 to Hemmi *et al.*, No. 5,378,715 issued 3 January 1995 to Stein *et al.*, and No. 5,382,569 issued 17 January 1995 to Cody *et al.* In addition, U.S. Patent No. 5,338,726 issued 16 August 1994 to Shinosaki *et al.* describes the chemical structure of ET converting
10 enzyme inhibitors. To date, however, antagonists of ET have not been approved for therapeutic use, although a number of investigators have postulated that ET antagonists could be used for conditions ranging from renal failure, endotoxic shock, asthma, angina, or diabetes to pulmonary hypertension and possibly other indications.

15 Under normal physiological conditions, ET can be found in almost all parts of circulation at very low levels. In general, in the normal rodent circulation ET is not found in elevated quantities and appears to have little detectable role in the normal regulation of vascular tone, i.e., there is no appreciable decrease in blood pressure when an ET antagonist is administered by injection in normal circulation. Further, at
20 present there does not appear to be any evidence suggesting that ET plays a physiological role even in a small portion of the circulation under normal circulatory conditions in experimental models. However, it is likely that the systemic circulation may appear to be normal when, in fact, specific regions of the circulation are undergoing pathophysiological changes such as occurs in conditions such as
25 erectile dysfunction (ED) (Adams *et al.*, *Int. J. Impot. Res.* 9:85-91, 1997).

Consequently, there are cardiovascular conditions which are traditionally treated in human beings by significant doses of NO or NO donors, such as glyceryl trinitrate (GTN) (0.2 mg/h and greater). However, these doses are known to induce systemic vasodilation and provoke considerable overall systemic side effects (The,

L.S. *et al.*, *Brit. J. Rheum.* 34:636, 1995). This is particularly so where a pathological condition exists only in certain major organs (e.g., heart, kidney, liver). As a result, a satisfactory method for promoting recovery of normal perfusion pressure in organs with certain pathologies without producing overall systemic hypotension has not been discovered.

Based on the understanding that a significant portion of underlying problems in clinical erectile dysfunction relates to "vascular" mechanisms, much of the current state-of-the-art research involves determining the contribution that the different vascular effector control systems make in normal and pathophysiological states. There is substantial understanding of hemodynamic events that lead to an erection, and yet the quantitative roles of each of the neuroeffector, humoral and local systems in these events remain poorly described. Since 1990, NO has been considered the primary non-adrenergic non-cholinergic neurotransmitter in the penis and has been presumed to be the primary mediator of corporal relaxation during erection (Ignarro L.J. *et al.*, *Biochem. Biophys. Res. Comm.*, 170:843, 1990).

It is well established that, for an erection to occur, neurally mediated (autonomic) vasodilation of the penile arterial blood vessel and the trabecular meshwork takes place (Lue, T.F. *et al.*, *J. Urol.* 137(5):829, 1987) permitting increased blood flow into the cavernous bodies of the penis. The expanding intra corporal volume compresses the effluent veins that lie between the erectile tissue and the surrounding fibrous, relatively inelastic, tunica albuginea. The outflow capacity is thereby decreased and entrapment of blood ensues, resulting in the transformation of the flaccid penis into its erect state (Lue, T.F. *et al.*, *J. Urol.* 137(5):829, 1987; Juenemann, K.P. *et al.*, *J. Urol.* 136(1):158, 1986; Lue, T.F. *et al.*, *J. Urol.* 130:1237, 1983; Weiss, H. *et al.*, *Ann Intern. Med.* 76:793, 1980). The level of arterial vascular tone (i.e., blood pressure) is one of critical importance in this process, although adequate perfusion pressure is also a necessary factor. The converse, detumescence, is mediated by the sympathetic nervous system (Saenz de Tajada, I. *et al.*, *Am. J. Physiol.* 254:H459, 1988; Juenemann, K.P. *et al.*, *Br. J. Urol.*

64, 1989).

The issue of "impotence" (defined as "a pattern of persistent or recurrent inability to develop or maintain an erection of sufficient rigidity for successful coitus") was discussed at a consensus conference of the National Institutes of Health (NIH) in Washington in December 1992 and has been clearly identified as having a wide range of causative or associated factors. The Massachusetts Male Aging Study (MMAS) has provided an updated view of the epidemiology of erectile dysfunction. It is accepted that the prevalence of impotence increases with age (Kinsey A.C. *et al.*, "Sexual Behaviour in the Human Male", W.B. Saunders: Philadelphia, 1948).

Severe or complete ED increases from 5 to 15% between 40 and 70 years of age, (Feldman, H.A. *et al.*, *J. Urol.* 151:54, 1994). ED has been shown to be "directly correlated with heart disease, hypertension, diabetes, associated medications, indices of anger and depression, and inversely with serum dehydroepiandrosterone, high density lipoprotein, cholesterol and an index of dominant personality."

It is now estimated that in North America there are more than 30,000,000 men with some form of ED, a significant increase from the figure of 10,000,000 quoted just 10 years ago (Shabsigh, R. *et al.*, *Urology* 32:83, 1988; Whitehead E., *Geriatrics* 43(2):114, 1988; Furlow, W.L. *et al.*, *Med. Aspects Human Sexuality* 19:13, 1985). From these figures it is also reasonable to estimate that as many as three million Canadian men may have a degree of ED. The direct cost of treating impotence is significant. Reliable figures from 1985 show that the cost of treating impotence exceeded 146 million dollars in the United States in that year alone (National Center for Health Statistics) and this number is just the estimated market size for one type of injectable therapy. The secondary effects and indirect costs associated with ED suggest that impotence and sexual dysfunction are medical icebergs. The consequences of sexual dysfunction may be seen in strains on the host relationship potentially leading to marital breakdown, violence, work related sequelae, deviant sexual behaviour and impacts on children, when present, that can carry the damage into a new generation of unwanted behaviours. If ED underlies even a small but

significant percentage of marital and family breakdown, then it adds vastly to the social and economic burden in society. The pragmatic issue is that large numbers of men are now being treated for ED and most of the treatments are fairly blunt instruments (e.g., intracavernosal injection (ICI)) of mixed vasoactive compounds, penile prosthesis insertion) with significant cost and complications (ICI: pain, priapism, dislike of the technique; prostheses: reoperation, infection, distortion of body image).

Administration of NO, or compounds which are able to deliver NO, have been suggested as possible therapies; however, these agents can provoke powerful yet inappropriate vasodilator responses (Brock *et al.*, *J. Urol.* 150:864, 1993). Such administration is often accompanied by a number of undesirable side effects related to systemic vasodilation which include headache, flushing, and hypotension. Consequently, there is a real need to provide methods whereby ED and other vascular disease may be quickly and effectively treated without any inappropriate side effects.

Summary of the Invention

Problems associated with localized imbalance in vascular tone, such as seen in ED, and which are hypothesized to be largely due to elevated ET, may be relieved by the administration of agents which are able to provide NO, by direct administration of NO, or by administration of an agent or agents which activates guanyl cyclase, such as, for example, YC-1, or other agents which prolong the actions of endogenous NO or of cGMP (a 2nd messenger molecule), such as phosphodiesterase (PDE) inhibitors, in minimal doses or microdoses, which heretofore had not been thought to result in effective treatment of an imbalance in vascular tone.

In the normal physiological state, there are sufficient quantities of NO present in the vasculature to maintain appropriate levels of ET (Banting *et al.*, *J. Hypertens.* 14:975, 1996). The addition of further NO has little impact on the effect of ET and

consequently any further vasodilation seen in such normal smooth muscle in the vasculature is likely attributable to NO's effects on the generation of cGMP, with cGMP resulting in decreased levels of Ca^{++} . In certain pathological conditions, such as diabetes and cardiovascular disease, and/or as a consequence of age, tissue is
5 unable to provide satisfactory levels of NO in order to suppress normal levels of ET present in smooth muscle tissues.

As such, physiological conditions where NO production is reduced in a specific local circulation, such as male ED, indicate that suppression of ET activity should offer an effective treatment. Consequently, in accordance with this, the
10 present invention provides for the use of an agent or agents which directly or indirectly generates NO at dosages which are about one half to about one twentieth ($1/2$ to $1/20$) of those known to induce vasodilation in "normal" circulations, and consequently exert almost no effect in the normal vasculature. As such these low doses, or "microdoses" are hypothesized to normalize the balance between NO and
15 ET. The range of about $1/2$ to about $1/20$ is derived from the observation that at doses which are below about $1/2$ the normal dose, systemic effects are generally no longer seen. At about $1/20$ the normal dose, however, the desired effect is also generally no longer observed, i.e., there is no effect.

According to one aspect of the present invention the concept of "low-dose" agents which directly or indirectly generate NO, or prolong the action of NO, or
20 enhance the cGMP 2nd messenger system, such as PDE inhibitors, is also applicable to any other peripheral pathological conditions where, regardless of the origin, NO is at least partially inhibited.

According to a further aspect of the present invention, there is provided a
25 method for restoring normal vascular tone in vasculature where NO levels are depleted and restoration of such levels may be achieved by addition of NO at levels which do not appreciably alter normal systemic vascular tone.

According to yet a further aspect of the present invention there is provided a method to treat any condition where NO levels are at least partially inhibited or

reduced, wherein the method comprises administration of NO or one or more NO donors, or one or more agents which activate guanyl cyclase, by oral, sublingual, buccal, intravenous, transdermal, or any other effective route, in concentrations that are about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature. Preferably, the concentrations of the method of the present invention are about one quarter to about one twentieth ($\frac{1}{4}$ to $\frac{1}{20}$) of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature. Still more preferably, the concentrations of the method of the present invention are about one eighth to about one sixteenth ($\frac{1}{8}$ to $\frac{1}{16}$) of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature.

According to one aspect of the present invention there is provided a method to treat instances of renal disease associated with excessive vasoconstriction where NO levels are at least partially inhibited, wherein the method comprises administration of NO or one or more NO donors, or one or more agents which activate guanyl cyclase, by oral, sublingual, buccal, intravenous, transdermal, or any other effective route, in concentrations that are about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in "healthy" or normal regions of the circulation such as the coronary or skeletal muscle vasculature.

According to a another aspect of the present invention there is provided a method to treat premature aging of the skin associated with inappropriate vasoconstriction of the skin vasculature which is associated with at least partial inhibition of NO levels in the skin, wherein the method comprises administration of NO or one or more NO donors, or one or more agents which activate guanyl cyclase, by oral, sublingual, buccal, intravenous, transdermal, or any other effective route, in concentrations that are about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature.

According to a further aspect of the present invention there is provided a method to treat male ED caused at least by partial inhibition of NO in the penile vasculature wherein the method comprises administration of NO or one or more NO donors, or one or more agents which activate guanyl cyclase, by oral, sublingual, buccal, intravenous, transdermal, or any other effective route, in concentrations that are about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature.

According to yet a further aspect of the present invention there is provided a method to treat ED comprising administering to a subject in need thereof a quantity of glyceryl trinitrate (GTN) by any route, for example, oral, sublingual, transdermal, intravenous, or inhalation, that provides plasma concentrations of below about 250 pg/ml of the GTN, so that ED is treated.

According to one aspect of the present invention there is provided a low-dose transdermal "patch" with short term release of, for example, GTN or any other effective provider of NO, such as, for example, one or more NO donors or one or more agents which activate guanyl cyclase, over a period of time less than 6 hours (as opposed to 12, 18 or 24 hour release), which restores normal vascular tone in an affected local vascular bed, such as the pudendal or penile vasculature of men with ED, without inappropriately affecting systemic vascular tone or causing hypotension, and preferably, but not necessarily, without inducing tolerance (Bennett *et al.*, *Circ. Res.* 63:693, 1988) to the effects and/or biotransformation of a NO releasing compound to its releasing form.

According to a further aspect of the present invention, there is provided a low dose or microdose "patch" with long term release of, for example, GTN, or any other effective provider of NO, such as, for example, one or more NO donors, or one or more agents which activate guanyl cyclase, over a period of time greater than 6 hours (typically 12 to 24 hour release), which restores normal vascular tone in an affected local vascular bed, such as the pudendal or penile vasculature of men with ED, without inappropriately affecting systemic vascular tone, and preferably, but not

necessarily, without inducing tolerance to the effects of a NO releasing compound and/or biotransformation of a NO releasing compound to its releasing form.

5 According to a further aspect of the present invention there is provided a method to treat female sexual dysfunction (SD) wherein the method comprises administration of NO or one or more NO donors, or one or more agents which activate guanyl cyclase, by oral, sublingual, buccal, intravenous, transdermal, or any other effective route, in concentrations that are about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in "healthy" or normal regions such as the coronary or skeletal muscle vasculature.

10 According to yet a further aspect of the present invention there is provided a method to treat SD comprising administering to a subject in need thereof a quantity of GTN by any route, for example, oral, sublingual, transdermal, intravenous, or inhalation, that provides plasma concentrations of below about 250 pg/ml of the GTN, so that SD is treated.

15 According to one aspect of the present invention there is provided a low-dose transdermal "patch" with short term release of, for example, GTN, or any other effective provider of NO such as, for example, one or more NO donors or one or more agents which activate guanyl cyclase, over a period of time less than 6 hours (as opposed to 12, 18 or 24 hour release), which restores normal vascular tone in an affected local vascular bed, such as the pudendal, cervical or vaginal vasculature of women with SD, without inappropriately affecting systemic vascular tone or causing hypotension, and preferably, but not necessarily, without inducing tolerance (Bennett *et al.*, *Circ. Res.* 63:693, 1988) to the effects and/or biotransformation of a NO releasing compound to its releasing form.

25 According to a further aspect of the present invention there is provided a low dose or microdose "patch" with long term release of, for example, GTN, or any other effective provider of NO, such as, for example, one or more NO donors or one or more agents which activate guanyl cyclase, over a period of time greater than 6 hours (typically 12 to 30 hour release), which restores normal vascular tone in an

affected local vascular bed such as the pudendal, cervical or vaginal vasculature of women with SD without inappropriately affecting systemic vascular tone, nor inducing tolerance to the effects of a NO releasing compound and/or biotransformation of a NO releasing compound to its releasing form.

5

Definitions

	GTN	glyceryl trinitrate
	ISMN	isosorbide 5-mononitrate
	ISDN	isosorbide dinitrate
10	PETN	pentaerythritol tetranitrate
	ETN	erythrityl tetranitrate
	SNP	sodium nitroprusside
	SIN-1	3-morpholiniosydnonimine molesidomine
	SNAP	S-nitroso-N-acetylpenicillamine
15	SNOG	S-nitrosoglutathione
	NOHA	N-hydroxy-L-arginine
	cAMP	cyclic adenosine monophosphate
	cGMP	cyclic guanosine monophosphate
	L-NAME	N ^o nitro-L-arginine methyl ester
20	IP ₃	inositol-1,4,5-triphosphate
	RIHP	renal interstitial hydrostatic pressure
	T	tumescence
	R	rigidity

25 "Applying various forms of NO" as used herein includes administering NO donor or NO producing agents.

"Enhancing penile erection" as used herein is understood to mean increasing physical size and improving tumescence and/or rigidity of a penis, preferably so that it is capable of intromission.

"Erection of good quality" and "effective erection" are used herein interchangeably to mean adequate for vaginal penetration (i.e., intromission, or intercourse).

5 "NO donor", "NO producing agent", "NO delivering compound", "NO generating agent" and "NO provider" are used interchangeably in this specification and include all compounds which donate NO through biotransformation, compounds which generate NO spontaneously, compounds which spontaneously release NO, or any other compounds which otherwise generate NO or an NO-like moiety and include: glyceryl trinitrate, isosorbide 5-mononitrate, isosorbide dinitrate, 10 pentaerythritol tetranitrate, erythrityl tetranitrate, sodium nitroprusside, 3-morpholinosydnonimine molsidomine, S-nitroso-N-acetylpenicillamine, S-nitrosoglutathione, N-hydroxy-L-arginine, S,S-dinitrosodithiol, or NO gas, or a functional equivalents thereof. In some cases, NO is generated by activation of guanyl cyclase.

15 "Penis" as used herein may be interpreted to apply equally to clitoris in so far as there is substantial equivalence between penile and clitoral erectile tissue. "Sexual dysfunction" (SD) as used herein includes aspects of female dysfunction and urogenital aging such as decreased vaginal lubrication, decreased vaginal engorgement, pain during intercourse such as, for example, dyspareunia, urogenital 20 infections; and urogenitalia as affected by post-menopause, diabetes, vascular disease, estrogen depletion conditions, sexual dysfunction, and idiosyncratic vaginal dryness, respectively.

"Various forms of NO" as used herein is understood to mean any one of NO[•], NO⁺ and NO⁻, preferably NO⁺ and NO⁻, and can include as an alternative CO (carbon 25 monoxide) in its various forms, which produces an equivalent effect to NO.

"Without inappropriately affecting systemic vascular tone" as used herein means not affecting mean arterial pressure so as to produce inappropriate systemic vasodilation with effects such as hypotension, headache, flushing.

Brief Description of the Drawings

Embodiments of the invention will now be described, by way of example, with reference to the accompanying drawings, wherein:

Figure 1 is a graphical representation of results from measurements of mean arterial blood pressure in rats and illustrates the differential impact on mean arterial pressure (mmHg) of infusions of sodium nitroprusside in control and L-NAME treated rats.

Figure 2 is a representative tracing of measured mean arterial pressure in a rat demonstrating the contrasting effect of low level infusions of sodium nitroprusside in control and acute L-NAME treated rats.

Figure 3 is a graphical representation of measurements of mean arterial blood pressure illustrating an enhanced glyceryl trinitrate dose response curve in rats treated with control and L-NAME, acutely.

Figure 4 is a schematic diagram depicting a range representative of a "normal" vascular controlling "window" of NO with comparisons of control and L-NAME treated rats.

Figure 5 is a representative tracing of changes in mean arterial pressure and renal interstitial hydrostatic pressure (RIHP) in respect of renal circulatory blockade of NO synthase.

Figure 6 is a representative tracing of changes in mean arterial pressure and renal interstitial hydrostatic pressure (RIHP) in respect of the capacity of L-arginine to induce decreases in RIHP in connection with the renal circulation.

Figure 7 is a representative tracing of changes in mean arterial pressure and renal interstitial hydrostatic pressure (RIHP) in respect of decreasing RIHP following removal of L-arginine in the renal circulation.

Figure 8 is a graph comparing the sensitivity of mean arterial pressure to the NO donor sodium nitroprusside (SNP, right) and to isopropyl norepinephrine (INE, left), in the presence and absence of the NO synthase blocker L-NAME.

Figure 9 is a cumulative dose response curve of mean arterial pressure to

prazosin, an α_1 blocker, in the presence and absence of NO synthase blockade (L-NAME).

Figure 10 is a graph illustrating a change in mean arterial pressure from control conditions in the presence of SNP low-dose, SNP + L-NAME, and L-NAME alone.

Figure 11 is a graph illustrating a dose-response relationship of mean arterial pressure for different doses of the nitric oxide donors SNP (top panel) and GTN (bottom panel), before and after both nitric oxide synthase blockade (L-NAME) and endothelin receptor antagonism (PD145065).

Detailed Description of the Invention

The physiological role of NO has been described as that of a chronic vasodilator agent acting to persistently "offset" the actions of a number of vasoconstriction systems. However, as discussed by Banting *et al.* (*J. Hypertens.* 14:975, 1996), based on this rationale, this would reveal that the NO vasodilator system would normally have an overall activity level at close to 70% of the NO system's maximal capacity, leaving the system little reserve or gain to protect against further deviations in mean arterial pressure and blood flow homeostasis.

Banting *et al.* (*J. Hypertens.* 14:975, 1996) proposed that the chronic role of NO, *in vivo*, is not that of a chronic vasodilator system but rather an inhibitor of the activity of local vasoconstrictor agents, such as ET. That is, the hypertension following blockade of NO production was completely reversed with the administration of an ET receptor antagonist. This finding, combined with the understanding that ET appears to play almost no role in the "normal" physiological control of resting mean arterial pressure, indicates that the function of NO may be different than previously proposed. The results of the **Quantitative Study** as detailed below, in summary, indicate that:

- (i) the amount of NO required to completely restore mean arterial pressure in L-NAME treated rats from hypertensive levels is much less

than that required to lower mean arterial pressure under control conditions;

- (ii) the increase in sensitivity to NO and NO donors allows for as little as 1/20th of the standard concentration to provide a given vasodilator response in NO synthase blockade rats;
- (iii) the level of NO required to completely reverse the hypertension in the chronic phase of L-NAME induced hypertension without altering blood pressure in control rats is at the same level as required to reverse the hypertension following acute NO synthase blockade; and
- (iv) the similar mean arterial pressure lowering, in both treated and control rats, that occurred with the administration of "high" concentrations of sodium nitroprusside suggests that signalling processes have been "normalized" at approximately 10-12 ug/kg per minute of sodium nitroprusside infusion.

Banting *et al.* (*J. Hypertens.* 14:975, 1996) have suggested that a major role of NO is not to function as a chronic vasodilator, but rather a chronic inhibitory regulator of ET-mediated vasoconstriction. In that study it was demonstrated that ET-mediated vasoconstriction does not contribute to the maintenance of vascular tone in a circulation with an intact NO generating system. Also, it was demonstrated that almost the entire L-NAME induced hypertension was both prevented and/or reversed via the administration of an ET receptor antagonist. The results detailed below under the heading **Quantitative Study** quantitate the exogenous level of NO required to restore this regulatory balance and, taken together, we propose that NO functions to suppress ET-mediated vasoconstriction within a physiological "window" as illustrated in Figure 4, in a manner which is unrelated to its direct role as a vasodilator substance. Indeed, it is proposed that the mechanism of action of the NO donors delivered at the micro doses disclosed herein, are not vasodilatory; — rather they act to suppress the endogenous role of endothelin.

These conclusions and their applicability to other microcirculations are

supported by studies conducted in the kidney in connection with measurements of renal interstitial hydrostatic pressure and mean arterial pressure as reported below under the heading: ***Effect of NO in Renal Vasculature.***

5 In a further application of the present invention, studies conducted in human males with a low-dose "patch" with up to 3 hours of release of effective NO, demonstrate that providing a "microdose" of NO restored normal erectile function. The results of these studies are detailed below under the heading: ***Clinical Data: For erectile dysfunction reversal with microdoses of NO.***

10 Accordingly, the present invention provides a method for restoring normal vascular tone through the use of an agent or agents which directly or indirectly generate(s) NO at dosages of about 1/2 to about 1/20 of those currently used in clinical applications.

15 The method of the present invention is also applicable to any other peripheral pathological conditions where regardless of the origin, NO is at least partially inhibited. In respect of the conditions detailed below evidence in the literature demonstrates that the production of NO is, at least partially, inhibited. This leads to an imbalance between NO and ET in favour of ET. The method of the present invention restores this balance with 'quantities' of NO that do not induce vasodilation, or not appreciably, in 'healthy' or non-NO deficient circulations.

20 Conditions wherein the present invention has application include any condition where regional circulation exhibits inappropriate vasoconstriction, such as, for example, ED, conditions associated with female sexual dysfunction involving vaginal and/or pelvic circulation, Raynaud's phenomenon (a condition of the fingers with constrained blood flow), as well as conditions associated with inappropriate

25 vasoconstriction in, for example, cutaneous and/or dermal (for example in reconstructive surgery, as well as scleroderma and diffuse cutaneous systemic sclerosis), cerebral, and renal circulations. Concerning ED, the therapy of the present invention can be used to restore "normal" vasculature response in a patient who has undergone a radical prostatectomy procedure where normal circulatory

control may not return for 6-8 months post operatively.

The therapy of the present invention therefore has application in female dysfunction and urogenital aging conditions such as vaginal lubrication, vaginal engorgement, pain during intercourse (dyspareunia), urogenital infections; and urogenitalia as affected by post-menopause, diabetes, vascular disease, estrogen depletion conditions, sexual dysfunction, and idiosyncratic vaginal dryness, respectively. In such situations, the therapy of the present invention works to improve blood flow to the vagina leading to vaginal engorgement promoting better circulation (lubrication).

It is understood that any agent which provides a dose range of NO is intended to be included within the scope of the present invention. Thus, in accordance with the invention, an agent may be NO, an NO-like substance, a substance which directly releases NO (e.g., an NO donor) or causes the release of NO, a substance having NO-like activity or effect, a substance which directly activates guanyl cyclase, or a substance that prolongs the actions of the 2nd messenger molecule cGMP, such as a PDE inhibitor. Regardless of which of such agent or agents is employed in the methods of the invention, what is important is that an enhanced level of NO is achieved, or the effect of an enhanced level of NO is achieved. The enhanced level is delivered to the vasculature such that normal vascular tone is achieved by a mechanism which involves normalizing levels of ET, thereby restoring the tone of the target microvasculature.

It should also be understood that the methods of the invention include the administration of an agent or agents as described above, (i.e, NO, an NO-like substance, a substance which directly releases NO (e.g., an NO donor) or causes the release of NO, a substance having NO-like activity or effect, or a substance which directly activates guanyl cyclase), either acutely or chronically. Acute administration is administration of a finite duration and can be, for example, as short as that associated with treatment of ED, wherein administration takes place for a prescribed period of time (e.g., up to several hours) at discrete instances when the

5 affect of the administration (penile erection) is desired. Acute administration can also be of longer, finite duration. Chronic administration is administration of a continuous and indefinite duration, and can be, for example, administration associated with a chronic condition such as that involving inappropriate vasoconstriction of the renal vasculature, where chronic administration is necessary to restore normal renal function.

10 The methods of the present invention are also applicable to treating vascular conditions wherein there is no appreciable decrease or inhibition of endogenous NO. Such vascular conditions occur, for example, when endothelial cells experience high shear from rapid movement of red blood cells passing in contact with, or in close proximity to, endothelial cells, resulting in increased ET. The methods of the invention restore NO/ET balance. High shear is associated with, for example, drug therapy and atherosclerosis.

15 While the invention has been particularly shown and described with reference to certain embodiments, it will be understood by those skilled in the art that various other changes in form and detail may be made without departing from the spirit and scope of the invention.

20 All scientific papers and patents referred to in this specification are incorporated in totality by reference herein.

Quantitative Study

25 The study detailed herein quantitatively characterizes the level of exogenous NO required to restore 'normal' vascular function following the acute and chronic blockade of NO synthase. Referring to the Figures, and in particular Figure 1, the concentration-response relationship with increasing levels of sodium nitroprusside (SNP) is markedly shifted for both acute and chronic NO synthase blockade treated rats. The concentration-mean arterial pressure response curves were divided into two groups, concentrations that did not and those that did induce a lowering of mean arterial pressure in the controls. A similar increased sensitivity to low levels of SNP

in the L-NAME treated rats was demonstrated both in acute and chronic phases of NO synthase blockade treatment. This level of SNP administration (0.5 to 8 ug/kg per minute) did not significantly decrease mean arterial pressure in control rats but did result in a marked decrease in mean arterial pressure.

5 Referring now to Figure 2 there may be seen an illustrative example of this finding: the depressor response to 2 ug/kg per minute in the NO synthase blockade phase was superimposed over the 2 ug/kg per minute response in the control period. The mean arterial pressure lowering induced by 12-32 ug/kg per minute induced a similar lowering in both control and treated rats (see Figure 1, lower panel),
10 indicating a convergence with respect to the depressor response to SNP.

Referring now to Figure 3 it may be seen that the cumulative glyceryl trinitrate concentration-mean arterial pressure response demonstrated results similar to SNP. The cumulative concentration-mean arterial pressure response curve was also shifted leftward in the NO synthase blockade treated rats. Again, the similar trend of
15 concentrations that do not lower mean arterial pressure in the controls, almost completely reversed the NO synthase blockade induced hypertension.

Further studies assessed the impact of different vasoactive systems on the relationship between nitric oxide and endothelin and, more specifically, the low dose effects of nitric oxide and the suppression of local levels of endothelin (ET), which
20 cause vasodilation and a normalization of vascular tone in focal regions of the circulation that are, at least in part, NO deficient. In these studies, L-NAME was administered to increase blood pressure, and two other vasodilator agents were employed: isopropylnorepinephrine (INE, a B_1/B_2 agonist, see Figure 8) and the α_1 blocker prazosin (see Figure 9). INE activates a cAMP mediated signal transduction
25 system, and prazosin decreases the activity of an IP_3 signal transduction system.

Figures 8 and 9 provide more detailed information on the mechanism underlying the increased sensitivity of mean arterial pressure to NO following NO synthase blockade. As can be seen from Figures 8 and 9, neither of these two vasodilator agents was able to normalize the excessive vasoconstriction that follows

NO blockade.

Specifically, the two curves on the left-hand side of Figure 8 clearly demonstrate that the dose response of mean arterial pressure to INE is identical both before and after nitric oxide synthase blockade, indicating that the signal mediated by another vasodilator, INE, is unchanged in the presence or absence of nitric oxide. It is widely known that the mechanism of action of INE is via the activation of cAMP mediated signal transduction. In contrast, the curves on the right-hand side of Figure 8 indicate that there is an increase in sensitivity of mean arterial pressure to the NO donor SNP following the blockade of NO synthase.

From a mechanistic standpoint, INE causes vasodilation by the activation of the cAMP signal transduction system in the vasculature, whereas SNP causes vasodilation by the activation of the cGMP signal transduction system or by direct effects of NO (or an equivalent). The absence of a shift in sensitivity of mean arterial pressure to INE after NO blockade indicates that the increased sensitivity to NO donors such as SNP is independent of the cAMP signal transduction system, and may be directly dependent on the cGMP signal transduction system.

Figure 9 illustrates that increased sensitivity to nitric oxide donors occurs following NO blockade, but does not occur following treatment with an α_1 receptor antagonist (prazosin). As prazosin is a second vasodilator agent, similar to INE, these agents thus present two examples of different vasodilator mechanisms that do not influence the relationship between nitric oxide and endothelin, which relationship is in accordance with the methods of the present invention.

Therefore, to summarize the data presented in Figures 8 and 9, these data present examples demonstrating the relationship between nitric oxide and endothelin and the effects of a low-dose NO therapy on normalizing the affected circulation with respect to endothelin and maintaining and/or restoring vascular tone. These data also demonstrate that the imbalance in vascular tone that follows NO synthase blockade is not due to angiotensin II (data not shown), the sympathetic nervous system, or vasopressin (for example, via the IP_3 system), but is specific to the cGMP

signal transduction system or direct effects of nitric oxide or its equivalent. The data further indicate that the addition of an α_1 receptor antagonist, in contrast to an endothelin receptor antagonist, does not restore a normal level of vascular sensitivity. This is in accordance with the invention in that decreased levels of NO (via L-NAME) result in a marked up-regulation of endothelin, such as may be explained by an uncoupling of cGMP signalling.

Figure 10 is similar in principle to Figure 2 and provides a further quantitation of the 'microdose' level of exogenous NO that maintains a normal level of vascular tone following nitric oxide synthase blockade with L-NAME. The graph of Figure 10 illustrates that administering a level of SNP that lowers blood pressure by 8 mmHg completely prevents the development of hypertension following even maximal NO synthase blockade (which is usually 40 mmHg). Thus, in accordance with the invention, it can be seen that a microdose of SNP causes only a minimum level of vasodilation, but completely prevents the excessive vasoconstriction which occurs following high level NO synthase blockade using L-NAME.

Further evidence supports that nitric oxide chronically suppresses the effects of endothelin. As shown in Figure 11, even normal levels of NO induce changes in the vasculature by countering the actions of endothelin. In Figure 11 (top panel), there can be seen an increase in the sensitivity of mean arterial pressure to SNP following NO synthase blockade, and a decrease in the sensitivity to SNP in the presence of PD145065, an endothelin receptor antagonist. The rightward shift in the curve for PD145065 (under control conditions) clearly indicates that the response to NO by itself has been severely blunted. The bottom panel of Figure 11 similarly indicates that the increased sensitivity to the NO donor GTN in response to treatment with L-NAME is abolished in the presence of PD145065. These results indicate that endothelin receptors are required to maintain the normal level of nitric oxide vasodilatory capacity under control conditions.

As would be apparent to a person of ordinary skill in the art, it is reasonable to use the rat as a model for the affected vascular systems discussed herein, such

as, for example the pudendal and penis vasculature, and to extend such studies to appropriate dosages and therapies for other subjects, such as humans. As is evidenced by Mordenti, "Man versus Beast: Pharmacokinetic Scaling in Mammals", *J. Pharm. Sci.* 75:1028-1040 (1986) and similar articles, dosage forms for animals, such as for example rats, can be and are widely used directly to establish dosage levels in human applications. One of the present inventors contributed to the development of a bioassay of erectile function in a rat model at least as early as 1991 (Heaton *et al.*, *J. Urol.* 145:1099-1102, 1991), and also helped demonstrate in comparative tests of erectile function in humans and rats, that the narrow effective dose window of an orally administered drug, apomorphine, is almost identical when suitably adjusted for the differences in body weight as taught by Mordenti, cited above (Heaton *et al.*, *Urology* 45:200-206, 1995).

Effect of NO in Renal Vasculature

In essence, the kidney functions as a reverse filter in that almost all of the contents of the blood are filtered into beginning "urine" of the kidney and the kidney specifically reabsorbs what it wants to retain later (in the nephrons and loop of Henle). A physical force, called the renal interstitial hydrostatic pressure (RIHP), in the tissues of the kidney can have a profound effect on this process. Increased RIHP presents a physical force opposing the reabsorption process affecting reabsorption of sodium, water and other parts of the beginning filtered urine. If not reabsorbed the result is a diuretic response (natriuresis). In cases of low RIHP the opposite occurs. Removal of the physical force opposing reabsorption results in volume accumulation (more blood volume = higher blood pressure). This pressure-mediated natriuresis is explained by the fact that some of the blood vessels in the kidney are not "auto regulated" (much like the penile vasculature), such that the pressure generated in the kidney tissue (RIHP) is dictated by the level of vasoconstriction/kidney perfusion. Excessive vasoconstriction results in a decrease in RIHP, thereby removing a strong mechanism by which the kidney prevents the

reabsorption of the contents of filtered urine.

It is possible to simulate a type of renal failure, where excessive vasoconstriction predominates. In these circumstances a volume conservation situation is created. This simulation is carried out by treatment of anaesthetized rats which are fitted with instrumentation for measuring RIHP and mean arterial blood pressure with NO synthase blockers (L-NAME). The results from such experiments are depicted in Figures 5, 6, and 7 which illustrate "raw" pressure tracings of an example of these preparations (Methods written below, see **EXPERIMENTAL ASPECTS OF STUDIES REPORTED: Effects of NO in Renal Vasculature**). The traces illustrate that administration of NO synthase blocker L-NAME, results in a sharp decline in RIHP (Figure 5) consequent upon vasoconstriction of the vasculature. Administration of the NO precursor L-Arginine (150-200 mg/kg per min) reversed the renal vasoconstriction (as measured by RIHP changes - see Figure 6) but did not alter the level of mean arterial pressure. When the L-Arginine administration is stopped (with continued administration of L-NAME, the RIHP once again sharply declines as illustrated in Figure 7.

Clinical Data: For erectile dysfunction reversal with microdoses of NO

General Methods. Measurements were made in a clinical erectile dysfunction (ED) laboratory in patients with previously diagnosed ED. These patients were being evaluated for the purpose of optimization of their intracavernous injection dosing. All men were assessed using the Queen's University Human Sexuality Group Protocol (Kingston, Ontario, Canada).

Case 1

The first patient was a man with total erectile dysfunction of 18 months. History of hypertension, myocardial infarct (x2), coronary artery by-pass graft surgery, cerebral vascular accident and peripheral vascular disease. He was prescribed 0.2 mg/hr nitro patch (GTN, Ciba-Geigy) with no effect on systemic blood pressure. The

therapy with 0.2 mg/hr nitro patch was on 2 occasions with 100% successful intercourse. He had angina on the second try. This result supports the view that the quantity of NO administered was working at a level significantly lower than quantities typically used.

5

Case 2:

This patient was a man with no response to intra cavernosal injection therapy with conventional drugs. 0.2 mg/hr nitro patch (Ciba-Geigy) followed by 10 mcg PGE1. The patient experienced the first erection in 4 years (predating radical prostate surgery).

10

The concentration of glyceryl trinitrate (the typical form of NO donor derived from Nitrospray or Nitropatches) has been reported in the literature for such delivery routes to be in the order of 200 to 400 pg/ml of plasma (Sun *et al.*, *J. Clin. Pharmacol.* 35:390, 1995). Based on the clinical case studies performed, it was noted that the beneficial effects of a 0.2 mg/h patch are observed after 20 to 30 minutes. This corresponds to a plasma concentration of 100 pg/ml. In respect of the application of the invention to ED therapy a preferred range of NO agent (such as GTN) is a steady state plasma concentration of about 50 to about 200 pg/ml. This range corresponds to the "window" referred to above as illustrated in Figure 4.

15

20

Support for the proposal that the mechanism of action of the NO donors delivered at microdoses disclosed herein, are not directly vasodilatory, can be accomplished by giving animals NOS blocker high doses to create hypertension. This would be followed by infusion of an endothelin antagonist at a concentration which would almost completely reverse the hypertension induced by the NOS blocker. Subsequently, an NO donor, or combination of NO donors is infused. The concentration response curve from this infusion should now be similar to that obtained in control animals. In other words, the doses of NO which will cause a lowering of pressure will be at higher values, namely 10 to 24 times higher than in an L-NAME blocked animal alone thereby showing that high doses of NO are

25

vasodilatory and lower ones are, on their own, not vasodilatory. Rather, they act to suppress ET.

While the routes of administration of the NO donors reported here include intracavernous (IC) injection of NO donor, or by nasal spray or by patch, the present invention includes administration by means of topical creams, pharmaceutically acceptable organic and inorganic carrier substances suitable for parenteral, enteral, intraurethral, vaginal application, or transmucosal application via, for example, the respiratory tract (e.g., by inhalation, such as through intranasal application), which do not deleteriously react with the active compounds.

Compositions of the invention are administered to subjects in a biologically compatible form suitable for pharmaceutical administration *in vivo*. By "biologically compatible form suitable for administration *in vivo*" is meant a form of the active compounds of the invention to be administered in which any toxic effects are outweighed by the therapeutic effects of the active compounds of the invention. The term subject is intended to include living organisms in which a response can be elicited, e.g., mammals. Examples of subjects include humans, dogs, cats, mice, rats, and transgenic species thereof.

Administration of a therapeutically active amount of the therapeutic compositions of the present invention is defined as an amount effective, at dosages and for periods of time necessary to achieve the desired result. For example, a therapeutically active amount of active compounds of the invention may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of an agent or combination of agents of the invention to elicit a desired response in the individual. Dosage regimens may be adjusted to provide the optimum therapeutic response. For example, several divided doses may be administered or the dose may be proportionally reduced as indicated by the exigencies of the therapeutic situation.

The active compounds (e.g., SNP) may be administered in a convenient manner such as by injection (subcutaneous, intravenous, intracavernous, etc.), oral

administration, inhalation, transdermal application, rectal administration, urethral administration, vaginal administration, or intracavernous introduction. Depending on the route of administration, the active compound or compounds may be coated in a material to protect the compound(s) from the action of enzymes, acids and other natural conditions which may inactivate the compound(s), or facilitate or enable delivery of said compound(s).

A SNP composition (or other NO donor) or as a separate agent can be administered to a subject in an appropriate carrier or diluent, co-administered with enzyme inhibitors or in an appropriate carrier such as liposomes. The term "pharmaceutically acceptable carrier" as used herein is intended to include diluents such as saline and aqueous buffer solutions and vehicles of solid, liquid or gas phase. To administer an agent or agents of the present invention by other than parenteral administration, it may be necessary to coat the active compound(s) of the invention with, or co-administer the agent or agents of the present invention with, a material to prevent its inactivation. Liposomes include water-in-oil-in-water emulsions as well as conventional liposomes (Strejan *et al.*, *J. Neuroimmunol.* 7:27 1984). The active compound(s) may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils and other solutions. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms, stability enhancers and compounds to preserve physical characteristics that are needed for appropriate delivery.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases, the composition must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The pharmaceutically acceptable carrier can be a solvent or

dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, and sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating active compound (e.g., SNP) or compounds in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filter sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of an active ingredient of the invention plus any additional desired ingredient from a previously sterile-filtered solution thereof.

When an active compound(s) is suitably protected, as described above, the composition may be orally administered, for example, with an inert diluent or an assimilable edible carrier. As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such medica and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the

active compound, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the therapeutic treatment of individuals.

EXPERIMENTAL ASPECTS OF STUDIES REPORTED

Quantitative Study

Animals

Male Sprague-Dawley rats (325 to 400 g) obtained from Charles River Laboratories (Montreal, Canada) were housed individually under conditions of 12-hour light/dark cycle, with room temperature at 22 to 24°C, and were provided with Purina rodent chow and tap water *ad libitum* for at least 2 days before any experiments were started.

Measurement of MAP and Short Acting Drug Administration

The surgical method was based on the technique of Thompson *et al.* (*Hypertens.* 20:809, 1992). In brief, rats were anaesthetized with ketamine/xylazine (70/5 mg/kg i.p.), and the descending aorta distal to the kidneys was catheterized with small bore Teflon tubing (0.012 in i.d., 30 gauge, Cole-Parmer, Laval, Quebec, Canada) inserted into vinyl tubing (0.02 in i.d., 0.060 in, 23 gauge). The inferior vena cava was also catheterized distal to the kidneys with small bore Teflon tubing

(0.012 in i.d., 30 gauge, Cole-Parmer). The catheters were filled with heparinized saline (10 IU/ml) and held in place by a small amount of cyanoacrylate glue at the puncture site. The catheters were tunnelled subcutaneously and exteriorized at the back of the neck and sutured in place. Two days after surgery, MAP could be recorded (MacLab DAS, ADInstruments, Milford, MA). After connection, an equilibration period of approximately 30 minutes allowed for the determination of the steady state level of MAP before any recording began. Baseline MAP was determined from readings averaged over 5 minutes, taken from each rat at 15 minute intervals for at least 1 hour prior to the start of any experiment.

Sodium nitroprusside and glyceryl trinitrate concentration - mean arterial pressure response curves following acute and chronic N^w-nitro-L-arginine methyl ester.

Rats were randomly assigned treatment with N^w-nitro-L-arginine methyl ester (L-NAME) for 30 minutes (100 mg/kg, intraperitoneally) or 12 days (100 mg/kg, in drinking water) or tap water. Two days prior to the day of the experiment rats were instrumented with catheters, as described above. Following baseline measurements of MAP, rats were given a infusions of sodium nitroprusside (SNP, 0.5 to 32 ug/kg per minute dissolved in 0.9% sterile saline) with a step-wise increase in concentration every two minutes. Rats were allowed 30 minutes to recover from the SNP administration. Rats were then given infusions of glyceryl trinitrate (GTN, 0.5 to 32 ug/kg per minute dissolved in 0.9% sterile saline) with a step-wise increase in concentration every two minutes.

Throughout all of these pharmacological manipulations, MAP and HR were recorded at a sampling rate of 100 Hz and the data was stored on a disk drive for later analysis.

Effects of NO in Renal Vasculature

Surgical preparation: Experiments were performed in male Sprague-Dawley rates, 250 to 330g, obtained from Charles River Laboratories (Montreal, Canada).

Food (Purina rat chow) and water were provided *ad libitum* throughout the study. Rats were acclimatized for at least a week prior to experimentation in group housing under conditions of a 12-hour light/dark cycle, with room temperature at 22 to 24°C. 3 to 4 days prior to each experimental day, rats were instrumented with an aortic catheter as described in detail previously (Thompson *et al.*, *Hypertens.* 20:809, 1992). In brief, the rats were anaesthetized with ketamine hydrochloride (70 mg/kg i.p. (Rogarsetic, Rogar/STB Inc., Montreal, PQ) and xylazine hydrochloride (5 mg/kg i.p. (Rompum, Haver Inc., Etobicoke, ON) and placed on a heating pad to maintain a constant body temperature of 37°C. Additional ketamine was given, as necessary, during surgery. Following an abdominal incision, the descending aorta distal to the kidneys was catheterized with small-bore Teflon tubing (i.d. = 0.012 in, o.d. = 0.030 in, 30 gauge, Cole-Parmer, Laval, Quebec, Canada), inserted into vinyl tubing (i.d. = 0.02 in, o.d. = 0.060 in, 23 gauge). The catheter was filled with heparinized saline (50 IU/ml, Grade 1-A Heparin Sodium salt, Sigma Chemical Co., St. Louis) and held in place by a small amount of cyanoacrylate tissue glue (Lepage Ltd., Brampton, ON). The catheter was then flushed with a small volume (0.1 - 0.3 ml) of heparinized saline (50 IU/ml) and left entirely inside the abdominal cavity until the day of the RIHP measurements. The abdominal incision was sutured with 6-0 silk braided thread (Ethicon Ltd., Peterborough, Ontario). The rats were allowed to recover for 2 to 5 days, during which they were housed individually under the same conditions as described above. Rats were re-weighed following the recovery period and rats that had lost >30g were considered to have not sufficiently recovered from the surgery and hence, were excluded from the study (Haskins, S.C., Postoperative care, In: Methods of Animal Experimentation, Vol. III Part A, eds. Gay *et al.* (Academic Press Inc., Orlando, 1986)).

Experimental Procedure: The RIHP measurements were carried out in rats anaesthetized with thiobutabarbital sodium (Inactin, 100 mg/kg i.p., Byk-Gulden, Constance, Germany). This anaesthetic is used most frequently by others for the study of RIHP and PN and, in preliminary studies, yielded similar results to those

studies compared to a Ketamine/Xylazine anaesthetic preparation.

Rats were placed on a heating pad to maintain a constant body temperature of 37°C and the abdominal incision was reopened. Hematocrits were determined by the microcapillary tube method from arterial blood samples (300 µl) obtained from the arterial catheter. Only rates in which the hematocrit was found to be between 40 and 45 percent were subsequently used. Rates with hematocrits out of this range were considered to be hemodynamically challenged due to the blood loss during surgery and, since reductions in hematocrit can compromise cardiovascular function, were not used in this study (Haskins, S.C., Postoperative care, In: *Methods of Animal Experimentation*, Vol. III Part A, eds. Gay *et al.* (Academic Press Inc., Orlando, 1986); Pirkle *et al.*, *Endocrin.* 110:7, 1982; Houttuin *et al.*, *Am J. Physiol.* 223:63, 1972).

The arterial catheter, inserted previously, was attached to a pressure transducer (model CDX3, Cobe), and the pulsatile signal recorded using a physiograph (Beckman, model R511 or MacLab, ADInstruments Pty Ltd., Castle Hill, Australia). Small sections of the aorta (<0.5 cm proximal to the right kidney), mesenteric and celiac arteries were isolated and silk ligatures were placed loosely around the vessels.

RIHP was measured by implanting a cannula into the lateral side of the left renal cortex (Garcia-Estañ *et al.*, *Am. J. Physiol.*, 256:F63, 1989). The RIHP catheter was constructed by inserting a small core of porous polyethylene matrix material (2 mm, 35 µm pore size, Bel-Art Products, Pequannoc, NJ) into the heat expanded end of a polyethylene catheter (PE-50, i.d. = 0.023 in, o.d. = 0.038 in x 10-20 cm) such that approximately one-third of the matrix extends beyond the tubing. This technique ensures that tissue does not block the catheter while allowing for measurement of hydrostatic pressure (Roman *et al.*, *Am. J. Physiol.* 248:F190, 1985; Ott *et al.*, *J. Appl. Physiol.* 38:937, 1975). A 3 mm deep hole was made in the lateral surface of the left kidney with an electrocautery needle (26 gauge) and current was passed through the needle for approximately 3 to 5 seconds. Bleeding was

completely stopped by applying soft pressure with a cotton swab. The kidney was kept dry before implanting the RIHP catheter. The RIHP catheter was inserted into the hole and then sealed to the surface of the kidney capsule with cyanoacrylate. The RIHP catheter was similarly attached to the physiograph via the pressure transducer. Lack of pulsatile pressure ensured that the catheter was not in a blood vessel. To further check the location of the catheter two procedures were performed: (i) a small volume (100-200 μ l) of saline (10 IU/ml heparin) was slowly infused in order to obtain a characteristic increase in RIHP (Ott *et al.*, 1975) and (ii) a RIHP response to renal vein occlusion (RVO) was observed (Garcia-Estañ *et al.*, *Am. J. Physiol.*, 256:F63, 1989). Rats that failed to show these characteristic responses (Figure 2-1) were excluded from the experiment.

After a 10 to 15 minute equilibration period, the RVO response from the RIHP catheter and a pulsatile recording from the arterial catheter were re-confirmed and a baseline MAP and RIHP were recorded. During the control periods of the experiment, the exposed abdominal cavity was kept moist by covering with a wet gauze patch and 0.9% saline. In each experiment, arterial pressure was manipulated systematically from low to high MAP by sequentially tightening the ligatures around the upper aorta, celiac artery and mesenteric artery. Pressure changes were initially maintained for approximately 5 to 7 minutes in order to ensure that both MAP and RIHP equilibrate to the new steady state. As it was determined that RIHP equilibrated in less than 3 minutes all data was collected within this time. The RVO-RIHP response was checked periodically after reaching different arterial pressures. This sequence of arterial pressure manipulations was repeated after a 10 to 15 minute control period at baseline MAP and RIHP. The position of the RIHP catheter was verified at the end of each experiment and was typically found in the corticomedullary junction.

Those skilled in the art will recognize, or be able to ascertain through routine experimentation, equivalents to the specific embodiments described herein. Such equivalents are considered to be within the scope of the invention and are covered

by the appended claims.

WE CLAIM:

1. A method of treatment, in an organism, of a vascular condition, comprising administration of at least one agent at a level which enhances NO and which does not appreciably alter normal systemic vascular tone in said organism.
2. The method of claim 1 wherein said vascular condition is associated with a reduction in NO production.
3. The method of claim 1 wherein said vascular condition is associated with an attenuation of NO effect.
4. The method of claim 1 wherein said administration is acute.
5. The method of claim 1 wherein said administration is chronic.
6. The method of claim 1 wherein said at least one agent is selected from the group consisting of NO, CO, NO donors, CO donors, activators of guanyl cyclase, PDE inhibitors, and substances which produce an effect equivalent to that of NO.
7. The method of claim 6, wherein said at least one agent is an NO donor selected from the group consisting of glyceryl trinitrate, isosorbide 5-mononitrate, isosorbide dinitrate, pentaerythritol tetranitrate, erythrityl tetranitrate, sodium nitroprusside, 3-morpholinosydnonimine molsidomine, S-nitroso-N-acetylpenicillamine, S-nitrosoglutathione, and N-hydroxy-L-arginine.
8. The method of claim 1 wherein said administration is by at least one route selected from the group consisting of oral, sublingual, buccal, intravenous,

vaginal, rectal, transdermal, inhalation, enteral, and parenteral.

- 5 9. The method of claim 1 wherein said level of administration of said at least one agent is a concentration that is about $\frac{1}{2}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in an anatomical site of said organism lacking said vascular condition.
- 10 10. The method of claim 1 wherein said level of administration of said at least one agent is a concentration that is about $\frac{1}{4}$ to about $\frac{1}{20}$ of a concentration required to induce vasodilation in an anatomical site of said organism lacking said vascular condition.
- 15 11. The method of claim 1 wherein said level of administration of said at least one agent is a concentration that is about $\frac{1}{8}$ to about $\frac{1}{16}$ of a concentration required to induce vasodilation in an anatomical site of said organism lacking said vascular condition.
- 20 12. The method of claim 1 wherein said vascular condition is associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle, and gastro-intestinal tract.
- 25 13. The method of claim 1 wherein said vascular condition is associated with ED.
14. The method of claim 1 wherein said vascular condition is associated with female SD.
15. The method of claim 1 wherein said at least one agent which enhances NO is GTN, and said level of administration is a plasma concentration below about 250 pg/ml of GTN.

16. The method of claim 15 wherein said GTN is delivered via at least one route selected from the group consisting of oral, sublingual, buccal, intravenous, vaginal, rectal, transdermal, inhalation, enteral, and parenteral, and said level of administration is a plasma concentration below about 250 pg/ml of GTN.
- 5 17. The method of claim 1 wherein said at least one agent is administered transdermally via a low-dose patch with short term release of an effective amount of said at least one agent over less than 6 hours.
- 10 18. The method of claim 1 wherein said at least one agent is administered transdermally via a low-dose patch with long term release of an effective amount of said at least one agent over about 6 to about 18 hours.
- 15 19. The method of claim 1 wherein said at least one agent is administered transdermally via a low-dose patch with long term release of an effective amount of said at least one agent over a period greater than 18 hours.
- 20 20. The method of claim 12 wherein said vascular condition associated with the skin is a condition selected from the group consisting of Raynaud's phenomenon, scleroderma, and diffuse cutaneous systemic sclerosis.
- 25 21. The method of claim 14 wherein said female SD condition is selected from the group consisting of vaginal lubrication, vaginal engorgement, pain during intercourse, dyspareunia, urogenital infections; and urogenitalia effects selected from the group consisting of post-menopause, diabetes, vascular disease, estrogen depletion conditions, sexual dysfunction, and idiosyncratic vaginal dryness.
22. The method of claim 13 wherein said male ED condition is due to a radical

prostatectomy procedure where normal circulatory does not return post-operatively.

- 5 23. The method of claim 1 wherein said vascular condition is associated with physiological insult.
24. The method of claim 1 wherein said vascular condition is transient.

1/8




FIGURE 1

2/8

FIGURE 2





FIGURE 3



3/8

FIGURE 4





FIGURE 5



4/8

FIGURE 6





FIGURE 7



5/8

FIGURE 8






FIGURE 9





6/8

FIGURE 10




7/8

FIGURE 11



8/8

FIGURE 12



PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/21, 31/295		(11) International Publication Number: WO 98/58633
A3		(43) International Publication Date: 30 December 1998 (30.12.98)
(21) International Application Number: PCT/CA98/00603		(81) Designated States: AU, BR, CA, JP, MX, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 22 June 1998 (22.06.98)		
(30) Priority Data: 60/050,491 23 June 1997 (23.06.97) US 60/086,750 27 May 1998 (27.05.98) US		
(71) Applicant (for all designated States except US): QUEEN'S UNIVERSITY AT KINGSTON [CA/CA]; Kingston, Ontario K7L 3N6 (CA).		
(72) Inventors; and (75) Inventors/Applicants (for US only): ADAMS, Michael, A. [CA/CA]; 341 Renda Street, Kingston, Ontario K7M 5X9 (CA). HEATON, Jeremy, P., W. [CA/CA]; R.R. #1, Kingston, Ontario K7L 4V1 (CA). BANTING, James, D. [CA/CA]; Unit 1, Apartment 201, 47 Van Order Drive, Kingston, Ontario K7M 1B6 (CA).		
(74) Agent: STEEG, Carol, Miernicki; Parteq Innovations, Queen's University, Kingston, Ontario K7L 3N6 (CA).		(88) Date of publication of the international search report: 12 August 1999 (12.08.99)
(54) Title: MICRODOSE THERAPY OF VASCULAR CONDITIONS BY NO DONORS		
(57) Abstract <p>Methods for treating vascular conditions associated with localized imbalance in vascular tone, which are hypothesized to be largely due to elevated endothelin (ET), are provided. The methods involve administration of nitric oxide (NO), agents which are able to provide NO, such as NO donors, agents which activate guanyl cyclase, such as YC-1, or agents which prolong the actions of endogenous NO or cyclic guanosine monophosphate (cGMP; a 2nd messenger molecule), such as phosphodiesterase (PDE) inhibitors. According to the invention, such agents are administered in minimal doses or microdoses by any route known in the art, so as to provide dosages which are about one half to about one twentieth (1/2 to 1/20) of those known to induce vasodilation in "normal" circulations. The low doses of these agents effectively alleviate vascular conditions associated with a reduction in NO production or an attenuation of NO effect, by restoring balance in vascular tone while exerting almost no systemic effect in normal vasculature.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LJ	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No
PCT/CA 98/00603

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/21 A61K31/295

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	STIEF C G ET AL: "Preliminary results with the nitric oxide donor linsidomine chlorhydrate in the treatment of human erectile dysfunction." JOURNAL OF UROLOGY, (1992 NOV) 148 (5) 1437-40. JOURNAL CODE: KC7. ISSN: 0022-5347., XP002090606 United States see "results" see abstract	1

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" documents which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "G" document member of the same patent family

Date of the actual completion of the international search

26 May 1999

Date of mailing of the international search report

7.06.99

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040. Tx. 31 851 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

A. Jakobs

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 98/00603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HELLSTROM, WAYNE J. G. (1) ET AL: "Penile erection in the primate: Induction with nitric-oxide donors." JOURNAL OF UROLOGY, (1994) VOL. 151, NO. 6, PP. 1723-1727. ISSN: 0022-5347., XP002090607 see page 1725, table on 2nd column: see "discussion" see abstract; figure 1 see page 29, line 3-19	1-13, 22-24
X	WO 95 05172 A (ZONAGEN INC) 23 February 1995 see abstract see page 7, line 14 - page 14, line 21 see page 29, paragraph 3-19 see page 30, line 6-23	1-24
X	US 5 059 603 A (RUBIN DAVID) 22 October 1991 see abstract; examples 1-4	1-24
X	KIRKEBY H.J. ET AL: "Role of the L-arginine/nitric oxide pathway in relaxation of isolated human penile cavernous tissue and circumflex veins." ACTA PHYSIOL. SCAND., (1993) 149/3 (385-392). ISSN: 0001-6772 CODEN: APSCAX, XP002090608 United Kingdom see abstract	1-6, 8-14, 22-24
X	HEATON J.P.W.: "Preliminary studies, in rabbit penile cavernosal tissue, of the role of synthetic nitrovasodilators in erectile response." CURR. THER. RES., CLIN. EXP., (1989) 45/2 (278-284). ISSN: 0011-393X CODEN: CTCEA, XP002090609 United States see abstract see page 281, paragraph 3 - page 282, paragraph 1; figure 2	1-19, 22-24
X	KUROKAWA, HIROSHI ET AL: "Portal and systemic hemodynamic responses to a very low dose of nitroglycerin in cirrhotic patients with portal hypertension" NIPPON IKA DAIGAKU ZASSHI (1995), 62(1), 2-12 CODEN: NIDZAJ;ISSN: 0048-0444, XP002090610 see abstract; figures 1-5; tables 2,3 see page 5, paragraph 3 - page 8, paragraph 2	1-12,15, 16,21

-/-

INTERNATIONAL SEARCH REPORT

Inter: nal Application No
PCT/CA 98/00603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
X	GARDINER, S. M. ET AL: "Regional and cardiac hemodynamic responses to glyceryl trinitrate, acetylcholine, bradykinin and endothelin-1 in conscious rats: effects of NG-nitro-L-arginine methyl ester" BR. J. PHARMACOL. (1990), 101(3). 632-9 CODEN: BJPCBM; ISSN: 0007-1188, XP002090611 see abstract; figures 1-3; tables 1-3 see page 634, column 1 - page 636, column 2	1-12.15, 16.23.24
X	ANDERSON D C JR ET AL: "Topical nitrate treatment of impotence." ANNALS OF PHARMACOTHERAPY, (1993 OCT) 27 (10) 1203-5. REF: 17 JOURNAL CODE: BBX. ISSN: 1060-0280., XP002090612 United States see the whole document	1-13. 15-19, 22-24
X	WO 96 27372 A (INT MEDICAL INNOVATIONS INC ;ALLEN MICHAEL P (US)) 12 September 1996 see page 7, line 1 - page 12, line 14; claims 1-24; examples 1-9	1-19. 21-24
X	RENGANATHAN R ET AL: "Comparison of transdermal nitroglycerin and intracavernous injection of papaverine in the treatment of erectile dysfunction in patients with spinal cord lesions." SPINAL CORD, (1997 FEB) 35 (2) 99-103. JOURNAL CODE: CKK. ISSN: 1362-4393., XP002090613 ENGLAND: United Kingdom see abstract see page 101, column 1, paragraph 6 - column 2, paragraph 6	1-13, 15-19, 22-24
X	AHMAD S: "Scleroderma and impotence: response to nitroglycerin applied to the fingers and penis 'letter!'." SOUTHERN MEDICAL JOURNAL, (1990 DEC) 83 (12) 1495. JOURNAL CODE: UVH. ISSN: 0038-4348., XP002090614 United States see the whole document	1-12, 15-20, 23,24

-/-

INTERNATIONAL SEARCH REPORT

Intern: at Application No

PCT/CA 98/00603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No
A	JEREMY J.Y. ET AL: "Effects of sildenafil, a type-5 cGMP phosphodiesterase inhibitor, and papaverine on cyclic GMP and cyclic AMP levels in the rabbit corpus cavernosum in vitro." BRITISH JOURNAL OF UROLOGY, (1997) 79/6 (958-963). REFS: 22 ISSN: 0007-1331 CODEN: BJURAN, XP002090615 United Kingdom see the whole document	1-24
A	MARTEL A.M. ET AL: "Sildenafil." DRUGS OF THE FUTURE, (1997) 22/2 (138-143). REFS: 17 ISSN: 0377-8282 CODEN: DRFUD4, XP002090616 Spain see the whole document	1-24
X	BERKOW R.; FLETCHER A. (EDS.): "MERCK MANUAL OF DIAGNOSIS AND THERAPY" 1987, MERCK SHARP & DOHME RESEARCH LABORATORIES, RAHWAY, N.J. XP002103703 67991 see page 400	1-12, 17-19, 21,23,24
X	TRIGO-ROCHA F ET AL: "Sodium nitroprusside: physiologic effects as a nitric oxide donor in three species." INTERNATIONAL JOURNAL OF IMPOTENCE RESEARCH, (1995 MAR) 7 (1) 49-56. JOURNAL CODE: BUX. ISSN: 0955-9930., XP002103698 ENGLAND: United Kingdom see abstract	1-7, 9-13,17, 22-24
X	VARGAS F ET AL: "VASCULAR REACTIVITY AND FLOW-PRESSURE CURVE IN ISOLATED KIDNEYS FROM RATS WITH N-NITRO-L-ARGININE METHYL ESTER-INDUCED HYPERTENSION" JOURNAL OF HYPERTENSION, vol. 14, no. 3, 1 March 1996, pages 373-379, XP000647927 see page 379, column 1, paragraph 2-3 see abstract; figure 3; example 2	1-12,23, 24
X	VARGAS, FELIX ET AL: "Endothelium-dependent and endothelium-independent vasodilation in hyperthyroid and hypothyroid rats" PHARMACOLOGY (1995), 51(5), 308-14 CODEN: PHMGBN;ISSN: 0031-7012, XP002103699 see abstract see page 312, column 1, paragraph 2 - page 313, column 2, paragraph 3	1-12,23, 24

-/-

INTERNATIONAL SEARCH REPORT

Intern 1al Application No

PCT/CA 98/00603

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
X	VARGAS F. ET AL: "Abnormal renal vascular reactivity to acetylcholine and nitroprusside in aging rats." GENERAL PHARMACOLOGY. (1997) 28/1 (133-137). REFS: 19 ISSN: 0306-3623 CODEN: GEPHDP, XP002103700 United States see abstract see page 134, column 2, paragraph 2-3; figures 1-3	1-12.23, 24
X	TARHAN, FATIH ET AL: "Effect of intracavernosal sodium nitroprusside in impotence" UROLOG. INT. (1996), 56(4), 211-214 CODEN: URINAC; ISSN: 0042-1138, XP002103701 see abstract see page 212, column 1, paragraph 4 - column 2, paragraph 4; tables 1.2	1-7,9-13
X	MCNAY J L: "PHARMACOLOGY OF THE RENAL CIRCULATION" AMERICAN JOURNAL OF MEDICINE, vol. 62, no. 4, 1 April 1977, pages 507-513, XP000561592 see abstract see page 508, column 2, paragraph 3-5	1-12.23, 24
X,P	KING R C ET AL: "Low - dose sodium nitroprusside reduces pulmonary reperfusion injury." ANNALS OF THORACIC SURGERY, (1997 MAY) 63 (5) 1398-404. JOURNAL CODE: 683. ISSN: 0003-4975., XP002103702 United States see abstract see page 1400, column 2, paragraph 2 - page 1403, column 2, paragraph 1; figures 1,3,6; table 1	1-11.23, 24
O,X	& KING R C ET AL: "Low - dose sodium nitroprusside reduces pulmonary reperfusion injury." FORTY-THIRD ANNUAL MEETING OF THE SOUTHERN THORACIC SURGICAL ASSOCIATION, 1996, see the whole document	1-11.23, 24
X	DE 43 05 881 C (LOHMANN THERAPIE SYST LTS) 3 March 1994 see claims 9,23	1-13, 17-19, 22-24

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 98/00603

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claim(s) 1-24
is(are) directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
1-14, 17-24 (partially) 15, 16
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☒ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-14,17-24 (partially), 15,16

Use of NO and/or glyceryl trinitrate in the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

2. Claims: 1-14,17-24 (partially)

Use of isosorbide 5-mononitrate and/or isosorbide dinitrate in the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

3. Claims: 1-14,17-24 (partially)

Use of pentaerythritol tetranitrate and/or erythrityl tetranitrate for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

4. Claims: 1-14,17-24 (partially)

Use of sodium nitroprusside for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

5. Claims: 1-14,17-24 (partially)

Use of 3-morpholinosydnonimine molsidomine for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

6. Claims: 1-14,17-24 (partially)

Use of S-nitroso-N-acetylpenicillamine for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

7. Claims: 1-14,17-24 (partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Use of S-nitrosoglutathione for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

8. Claims: 1-14,17-24 (partially)

Use of N-hydroxy-L-arginine for the treatment of diseases associated with an anatomical site selected from the group consisting of kidney, skin, penis, vagina, brain, skeletal muscle and gastro-intestinal tract.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

In view of the large number of compounds, which are defined by the general definition(s)/formulae used in claims 1-24 the search had to be restricted for economic reasons. The search was limited to the compounds for which pharmacological data was given and / or the compounds mentioned in the claims, and to the general idea underlying the application. (see Guidelines, chapter III, paragraph 2.3)

The requirements of Article 6 PCT are not met by claim 6 (partially) because the use of CO or CO donors are not fully supported by the description, not mentioned in the description.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter national Application No

PCT/CA 98/00603

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9505172 A	23-02-1995	AT 174795 T AU 696815 B AU 7523894 A AU 9716898 A BR 9407250 A CA 2169071 A CN 1128950 A DE 69415535 D EP 0714300 A ES 2127409 T JP 9501677 T NO 960549 A NZ 271567 A US 5565466 A ZA 9406123 A	15-01-1999 17-09-1998 14-03-1995 04-03-1999 24-09-1996 23-02-1995 14-08-1996 04-02-1999 05-06-1996 16-04-1999 18-02-1997 12-04-1996 19-12-1997 15-10-1996 20-03-1995
US 5059603 A	22-10-1991	NONE	
WO 9627372 A	12-09-1996	US 5698589 A AU 701328 B AU 5302596 A BR 9607974 A CA 2214418 A CN 1177294 A CZ 9702818 A EP 0814800 A HU 9801234 A JP 11501629 T NO 974075 A PL 322110 A SI 9620038 A SK 117597 A	16-12-1997 28-01-1999 23-09-1996 13-01-1998 12-09-1996 25-03-1998 17-12-1997 07-01-1998 28-01-1999 09-02-1999 27-10-1997 05-01-1998 30-04-1998 14-01-1997
DE 4305881 C	03-03-1994	WO 9418966 A	01-09-1994

THIS PAGE BLANK (USPTO)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)